

SUPRACONDYLAR FRACTURE FEMUR TREATED WITH INTRAMEDULLARY NAIL: A PROSPECTIVE STUDY

D. Venkateswara Rao¹, G. Chaitanya², Anvesh Sangepu³

HOW TO CITE THIS ARTICLE:

D. Venkateswara Rao, G. Chaitanya, Anvesh Sangepu. "Supracondylar Fracture Femur Treated with Intramedullary Nail: A Prospective Study". *Journal of Evolution of Medical and Dental Sciences* 2015; Vol. 4, Issue 65, August 13; Page: 11352-11361, DOI: 10.14260/jemds/2015/1637

ABSTRACT: BACKGROUND: Supracondylar fractures are one of the commonest fractures encountered in high velocity trauma which are associated with high morbidity and mortality.¹ Isolated fracture can itself lead to complications such as ARDS and pulmonary embolism¹. This necessitates early stabilization of fractures. Internal fixation is the choice of treatment in supracondylar fractures (AO type-A).² Retrograde supracondylar nail has shown to give one of the best results in terms of recovery, fracture union, return to work and the functional outcome. **METHODS:** 20 patients with supracondylar fracture femur were studied (AO type-A). Supracondylar fractures femur were treated by closed or open reduction and internal fixation by Retrograde supracondylar nail inserted through inter-condylar notch between July 2012 to September 2014 at our institution. The patients were evaluated clinically and radiologically for outcomes. All patients were followed up for an average of 12 months. Outcome was assessed using NEER'S SCORE.³ **RESULTS:** Supracondylar femoral extra-articular fractures in 20 patients were treated in this study with retrograde intramedullary supracondylar nail after closed or open reduction. The AO-ASIF fracture classification was used. All extra-articular fractures were selected for study. Six were compound fractures according to the Gustilo-Anderson classification,⁴ the fracture was stabilized with the chosen system at an average of 6.9 days post-trauma, with an average operative time of 65.9 minutes. In 15 cases closed reduction was possible while 5 required open reduction. Post-operatively all patients were shifted to continuous passive mobilization with early toe touch walking and gradually progressive weight bearing with appearance of clinical and radiological signs of union. Evaluation was done according to Neer's rating system.³ 65% excellent results were found. In the present study, road traffic accidents were observed to be the predominant cause of distal third femoral fractures in young patients. All fractures had a sound clinical and radiological union with an average radiological union time of 16.2 weeks and average full weight bearing time was 13.4 weeks. Average knee flexion was 108 degrees with an extensor lag of 4.15 degrees.

KEYWORDS: Supracondylar femur nail.

INTRODUCTION: In the few decades, rapid industrialization and the fast pace of life have brought both comforts and catastrophe like road traffic accidents and crippling many young lives. Supracondylar and intercondylar femoral fractures are often difficult to treat and they are notorious for many complications. In the early 1960s, there was a great reluctance towards operative management of this fracture because of high incidence of infection, non-union, malunion, inadequate fixation and lack of proper instruments, implant as well as antibiotics. Then, the traditional management of displaced fracture of supracondylar femur was along the principle of Watson Jones¹ & John Charnely.⁵ This comprised of skeletal traction, manipulation of fracture and external immobilization in the form of casts and cast bracings.⁵ These methods however, met with problems

ORIGINAL ARTICLE

like deformity, shortening, prolonged bed rest, knee stiffness, angulation, joint incongruity, malunion, quadriceps wasting, knee instability and post-traumatic osteoarthritis.

The trend of open reduction and internal fixation has become evident in the recent years with good results being obtained with the AO blade plate, dynamic condylar screw and other implant systems like intramedullary supracondylar nails. Supracondylar fractures tend to collapse into varus, due to strong adductors. During application of AO blade plate or dynamic condylar screw, the shaft of femur is often pulled laterally displacing the line of weight bearing, lateral to the anatomic axis of femur condyle. This creates rotational movements at the fracture site that causes pulling off the blade plate or condylar screws leading to fatigue fracture of the plates. Also, the presence of osteoporotic bone leads to fixation failures with screws and plates cutting the soft bone. In contrary, a retrograde intramedullary supracondylar nail has got distinct advantages of decreased blood loss, minimal soft tissue dissection, less operative time and reduced rate of infection.⁶

The purpose of this study is to evaluate the results of type supracondylar fracture of femur, treated by close/open reduction and internal fixation using retrograde intramedullary supracondylar nail.

AIMS & OBJECTIVES:

AIM: To study the management of supracondylar fractures femur by Retrograde Supracondylar Interlocking Nail at Government General Hospital Vijayawada attached to Siddhartha Medical College Vijayawada, between July 2012 to September 2014.

OBJECTIVES:

1. To study and evaluate the results of the retrograde nailing in supracondylar femoral fractures in relation to knee movements, early mobilization of patients and early weight bearing.
2. To study the role of supracondylar nail in avoiding post-operative varus/valgus deformities and shortening in supracondylar fractures femur.
3. To study the postoperative rehabilitation of retrograde nailing in supracondylar femoral fractures.
4. Compare the results with similar other studies.

MATERIALS & METHODS: In this study 20 patients with supracondylar fracture of femur without intercondylar extension were studied. All the cases treated in Government General Hospital, Vijayawada attached to Siddhartha Medical College between the periods of July 2012 to September 2014. The method used for fracture fixation was closed or open reduction and internal fixation with retrograde intramedullary supracondylar nail. The duration of follow up ranged from 4 months to 24 months. All the fractures in this series were post-traumatic. No pathological fracture was included in the study. Also supracondylar fractures in children were not considered. The study was restricted to fractures occurring at the region 9 cm proximal to lower end of the femur. Supracondylar fractures treated conservatively and fixed with other fixation systems like dynamic condylar screw, AO blade plate and condylar buttress plate are not included.

The following Protocol was Observed for Patients with Supracondylar Fractures of Femur on arrival:

1. General and systemic examination as well as local examination of the patient.

ORIGINAL ARTICLE

2. Thorough assessment of patient to rule out head/chest/abdominal/spinal or pelvic injury.
3. Evaluation of patient in terms of: age, sex, mode of trauma, period between injury and arrival.
 - Musculo-skeletal examination of patient to rule out associated fractures.
 - Stabilization of patient in terms of Airway, breathing and circulation by oxygen, transfusion of crystalloids and colloids as and when required.
 - Careful assessment of injured limb as regards to neurovascular status.
 - Primary immobilization of involved limb in Thomas splint.
 - Radiological assessment: Anteroposterior and true lateral views of injured limb including complete knee joint and distal femur.
 - Upper tibial skeletal pin traction with a Steinmann pin drilled under local anaesthesia followed by continuous traction given over Bohler-Braun splint.
 - Compound injuries were taken for cleaning and debridement under anaesthesia at the earliest with meticulous debridement. Wound swab sent for culture and sensitivity for proper antibiotic coverage. Fixation was delayed in all cases.
 - Injection ATS 1500 IU, Injection AGGS 30, 000 IU, broad spectrum injectable antibiotics and analgesics were administered for compound injuries as and when required.
 - Patient Selection:
 - Patients attended to Government General Hospital, Vijayawada with supracondylar fractures of femur with:

Inclusion Criteria:

1. Type-A fractures (AO classification).
2. Grade 1, 2 and 3A fractures (Gustilio-Anderson classification).
3. Medically fit patients.
4. Patients of both sex.
5. Patients in the age group of 18-75 years.

Exclusion Criteria:

1. Patients with type B and C fractures (AO classification).
2. Grade 3B fractures (Gustilio-Anderson classification).
3. Medically unfit patients.
4. Patients below 18 years and above 75 years.
5. Patients with pathological distal femoral fractures other than osteoporosis.
6. Patients lost in follow-up.
7. Distal femoral fractures with neurovascular compromise.

Implant Used:

- The implant used was supracondylar nail system with instrumentation set.
- The nails are available with outer diameter of 10, 11 and 12mm.
- The distal end is expanded to outer diameter of 13mm.
- The nails are available in lengths of 150, 200 and 250mm.
- There is 5 degree anterior bend and anterior bow for anatomic fit.

ORIGINAL ARTICLE

All sized nails have five interlocking holes in all lengths two proximal holes and three distal holes, which accept interlocking screws of 4.9mm thread diameter. The interlocking holes are medio-laterally directed.



Intramedullary Supracondylar Nail with Instrumentation

OPERATIVE PROCEDURE: The limb was scrubbed for 5 minutes with surgical betadine scrub followed by painting with povidone iodine and medicated spirit and draping with sterile drapes so that knee joint and distal thigh were in the operative field. A midline incision of 4cm was taken from inferior pole of patella up to tibial tuberosity. The paratenon over patellar tendon was sharply incised and patellar tendon was split in the midline along the direction of its fibers. A straight bone awl was inserted into the joint through the split tendon and positioned against the inter-condylar notch. The femoral attachment of posterior Cruciate Ligament is palpated and bone awl is kept just anterior to the Posterior Cruciate Ligament attachment.

The position of bone awl was checked under image intensifier in antero-posterior and lateral position. An entry point was made. The bone awl was then removed and guide wire passed through entry point. The bone awl was then removed and guide wire passed through entry point. The fracture was reduced under image intensifier control and guide wire passed in proximal fragment. The distal fragment was then reamed with flexible shafts. The predetermined nail of adequate diameter and length was then loaded over the jig with the help of conical bolt keeping in mind the side to be operated so that jig was placed laterally and convexity of nail facing anteriorly. The nail was then inserted over the guide wire through the entry point made previously through distal fragment and then proximal fragment.

Its position was confirmed on image intensifier and then depending on the length of the nail, the proximal holes were locked with the help of corresponding markings on the jig. After taking stab incision over the corresponding lateral skin, the soft tissues were separated by blunt dissection with the help of haemostat and drill sleeve and drill guide for 4.5mm drill bit were inserted through the fenestrations provided over the jig, through the stab incision flush with the lateral cortex. The lateral and medial cortex were drilled with 4.5mm drill bit.

Continuity of drill holes in both the cortices with the locking hole of nails was confirmed with sounding technique. The required length of locking bolt was measured with the help of depth gauge and self-tapping interlocking bolt of 4.9mm thread diameter passed from lateral to medial cortex engaging the locking hole in the nail.


ORIGINAL ARTICLE

Either single or both holes were locked proximally. Similarly, the distal holes were locked in one, two or three numbers using 4.9mm interlocking bolt. The jig was then disengaged, the joint was washed thoroughly to remove the debris, hemostasis achieved and incision closed in layers. Particular attention was paid to repair paratenon of patellar tendon.

Disadvantages:

1. Need for an arthrotomy if the hardware needs removal.
2. Patello femoral arthrosis.
3. Knee stiffness.
4. Nail migration into joint if not locked.
5. Cruciate ligament injury.


RESULTS: In this study 20 patients with supracondylar fracture of femur without intercondylar extension were studied. All the cases were treated in Government General Hospital attached to Siddartha Medical College Vijayawada between July 2012 to September 2014. The method used for fracture fixation was closed or open reduction and internal fixation with retrograde intramedullary supracondylar nail. The duration of follow up ranged from 4 months to 24 months. 85% good to excellent result were obtained using Neer's evaluation scoring system. In this study, the youngest case was 30 years old male and the oldest was 75 years. Overall mean age was 52.15 years. In males, it was 44.89 years and females it was 47.5 years.




Number of cases Male & Female

In this study left side affection was seen more than right side. 75% fractures were sustained due to road traffic accidents. 15% were fall from height. 2% was accidental fall. In the present study, there were 6 compound fractures, 4 being grade-II and 1 being grade-III according to Gustilo-Anderson's classification.

ORIGINAL ARTICLE



Average Operative time was 86.5 minutes. It is 82.3 minutes for type A1, 93 minutes for type A2 and 130 minutes for type A3 fractures. In majority of the fracture type of A1 and A2 were closed reduction was possible, while in A3 subtype, cases required open reduction. Maximum closed reduced fractures required less than or 90 minutes for operation. While maximum open reduction fractures required more than 90 minutes operative time. Average operative time for closed reduced fractures was 79.3 minutes. Average operative time for open reduced fractures was 108 minutes. 12 out of 15 cases done within 7 days were reduced closely. Whereas 4 out 5 cases done after 7 days needed open reduction. Average radiological union time was 17.3 weeks.




Average flexion attained in this study was 93.25 degrees. More than 110 degrees flexion range was observed in 50% of the cases. Average extensor lag in this study was 5.75 degrees. Out of 20 patients, one had shortening of 23mm. Local symptoms at distal screws was found to be the commonest complications like pain and loosening of screws. Long-term results were rated using the Neer's rating system, which allots points for pain, function, working ability, joint movements, gross

ORIGINAL ARTICLE

and radiological appearance. In 65% cases, there was good to excellent results. No statistically significant correlation was found between knee flexion and age. One case with 75 degree flexion in A1 type of supracondylar fracture had comminuted proximal tibial fracture, which delayed the mobilization and weight bearing. Seven of Ten cases with A2 type had >110 degree flexion (70%). 4 of 9 cases with A1 type had >110 degree flexion (44.5%). One case of A3 fracture had <90 degree flexion. 46.6% patients with closed reduction had >110 degree flexion while 80% patients with open reduction had >110 degree flexion.

Final knee flexion was better, the earlier the patient was operated. Average weight bearing for A1 type fractures was 8.84 weeks, Type A2 was 10.53 weeks and 8 weeks for type A3 fractures. 70% of cases done by closed method and 90% done by open method had full weight bearing within 8 to 12 weeks. Average weight bearing for closed reduction was 12.2 weeks, and for open reduction was 13 weeks.



Pre-operative X-ray



Post-operative X-ray

ORIGINAL ARTICLE

DISCUSSION: Mean age group reported in this study was 52.15 years. Gellman RE et al.⁷ (1996) reported 50 years as mean age group. Watanabe Y⁸ (2002) reported 64 yrs as mean age group. Our study coincides with those of Gellman R et al.⁷

Lucas SE⁹ (1993) with 39 mean age group, sex incidence was 13 males and 11 females. Gellman RE⁷ (1996) observed 10 males (mean age 39yrs) and 12 females (mean age 60yrs) with overall mean age of 50 yrs. Watanabe Y⁸ (2002) study, whose mean age was 64 yrs, with 4 male and 20 female patients. In the present study, there were 15 male patients with average age group 48.5yrs, and 5 female with average age of 63 years. Thus, in the study conducted by Gellman⁷ (1996) and Watanabe⁸ (2002), where female predominance was seen. The age group under consideration was older in contrast to the study by Lucas SE⁹ (1993) and the present study, where male predominance was observed.

Studies conducted by Schatzker et al.¹⁰ (1974), Yang RS et al.¹¹ (1990) and Leung KS et al.¹² (1991), demonstrated road traffic accidents as major causal factor. Lucas SE⁹ (1993) reported 79% RTA, 17% fall and 4% gunshot wound. In the present study, RTA accounted for 75% of cases and 15% resulting from fall from height and another 10% accounted for accidental fall. We also agree with the results of Gellmann Re⁷ (1996) who stated that high energy fractures occurred more in young, male patients and low energy falls caused fractures in older age group.

High percentage of compound injuries were found in the studies by Lucas SE et al.⁹ (1993), Iannacone et al.¹³ (1994). Lucas SE (1993) did not report any difference in the final outcome in terms of knee flexion and nature of fracture. In the present study, there was 6 compound fractures, one being grade I and 4 being grade II and one grade III according to Gustilo-Anderson classification⁴. Of the 6 cases, 5 were male and 1 was female. Among them 5 were due to RTA and one by accidental fall. Patients with Gustilo-Anderson⁴ grade II and III were operated as early as possible. The final outcome in terms of knee flexion was not affected by nature of fractures. 3 out of 6 compound injured had knee flexion of >110 degree (50%), while 7 out of 14 closed fractures had knee flexion >110 degree (55%). Extensor lag was found to be more in compound fracture with 13 out of 13 patients having extensor lag >10 degree as compared to 4 out of 13 for closed fractures.

In the present study, 6 patients had associated injuries. Of the 6 injured patients, 2 with ipsilateral proximal tibial fracture had knee flexion of average 82.5 degrees and extensor lag of >10 degree. Patient with patellar fracture had knee flexion of 110 degrees and no extensor lag. 2 patients with tibial spine fracture had average knee flexion of 102.5 degree and extensor lag of 9 degree. Thus, it appears that though significantly less number of patients in the present study had associated trauma, it seems to affect the final outcome. This can be attributed to delayed mobilization and delayed weight bearing in these patients.

Lucas SE⁹ (1993) observed that average injury-surgery interval of 6 days. Watanabe Y⁸ (2002) observed the average injury-surgery interval of 3 days. In the present study, the injury-surgery interval was 6.9 days. The interval between injury and surgery could be attributed to:

- Days lost by the patient in transit from periphery to the institution, which caters to the tertiary care needs.
- Medical fitness of the patient for necessary anaesthesia.
- Arrangement for funds consumed considerable time.

The injury-surgery interval and final knee flexion were found to be inversely proportional with 61.5% of fractures fixed within 7 days (13 cases) had >110 degree flexion as compared to

ORIGINAL ARTICLE

42.95% after 7 days (7 cases). This relationship could be explained on the fact that surgery interval affected the type of reduction with 13 out of 17 fractures operated within 7 days could be closed reduced. And closed reduction was directly proportional to final knee flexion. Average operative time in this study was 65.9 minutes. Majority of type A1 and A2 fractures required operative time >90 minutes and type A3 fractures required more than 90 minutes. Average radiological union in this study was 17.3 weeks. The patients which required more time for union were having some associated injuries which delayed the period of mobilization and partial weight bearing. In the present study, average knee flexion was 105 degree, which coincides with the previous other studies. Average extensor lag in this study was 5.4 degree.

In the present study, we had 2 cases of superficial infection, treated by antibiotics and debridement. 3 cases of distal screw related problems were treated by screw removal. One case of delayed union was treated by delaying full weight bearing. Anterior knee pain due to impingement might be attributed to faulty nailing technique. Improvement in the nailing technique can reduce the incidence of impingement.

CONCLUSIONS:

1. Retrograde intramedullary supracondylar nail is a good fixation system for distal third femoral fractures, particularly extra-articular type.
2. The operative time is lessened with decrease in blood loss, if closed reduction can be achieved by not disturbing fracture hematoma and soft tissue.
3. Even with open reduction, there is less soft tissue trauma and less post-operative stiffness.
4. Distal screw related local symptoms is a common problem and is related to implant technique and has a definite learning curve.
5. Utmost care require to avoid infection.
6. There is no non-union, less delayed unions and rates of angular or rotational malunions.
7. Non-requirement of bone graft decreases the morbidity associated with donor site.
8. Early surgery, closed reduction, at least 2 screws in each fragment and early post-operative knee mobilization are essential for good union and good knee range of motion, weight bearing and early return to work.
9. Thus, supracondylar nail is the optimal tool for many supracondylar fractures of femur. It provides rigid fixation in a region of femur, where a widening canal, thin cortices and frequently poor bone stock make fixation difficult. Surgical exposure for nail placement requires significantly less periosteal stripping and soft issue dissection than that of lateral fixation devices. Orthopaedic surgeons experienced with intramedullary nail will find the supracondylar nail a useful technique, but requires attention to prevent complications.

REFERENCES:

1. Wilson JN. Watson Jone's Fractures and Joint injuries. 6th ed., pg. 1003-1070 (1982).
2. Ruedi TP, Murphy WM. AO Principles of fracture management. Stuttgart, New York: Thieme (2001).
3. Neer CS, Gratham SA, Shelton ML et al Supracondylar fractures of adult femur. JBJs, Vol. 49-A, pg. 591-613 (1967).
4. Gustilo RB, Anderson JT. Prevention of infection in the treatment of 1025 open fractures of long bone. Retrospective and prospective analysis. JBJs, Vol. 58-A, pg. 453-58, June (1976).

ORIGINAL ARTICLE

5. Charnely John. The closed treatment of common fractures. 3rd ed., pg. 197-204
6. Leung KS, Shen WY, Mui LT. Interlocking intramedullary nailing for supracondylar and intercondylar fractures of distal part of femur. JBJS, Vol. 73-A, pg. 332-40 (Mar 1991).
7. Gellman RE, Guy D paiement, Hellary D Green Treatment of supracondylar femoral fractures with a retrograde intramedullary nail. CORR, No. 332: 90-97 (1996).
8. Ingman AM. Retrograde intramedullary nailing of supracondylar femoral fractures: Design & Development of a new implant. Injury, 33(8); 707-12: (2002 Oct).
9. Lucas. Quoted by Rockwood CA, Green DP. Fractures in adult, 45th ed, Vol. II, pg. 1972-93 (1996).
10. Schatzker J, Horne G, Wadell J. The Toronto experience with supracondylar fractures of femur. Injury, 6, pg. 113-28 (1975).
11. Yang-Rong-Sen, Hwa-Chang Liv et al. Supracondylar fractures of the femur. J Trauma, Vol. 30, pg. 315-319 (Mar 1990).
12. Leung KS, Shen WY, Mui LT. Interlocking intramedullary nailing for supracondylar and intercondylar fractures of distal part of femur. JBJS, Vol. 73-A, pg. 332-40 (Mar 1991).
13. Iannacone WM, Bennett FS, Delong WG Initial experience with the treatment of supracondylar femoral fractures using the supracondylar nail: A preliminary report. J Ortho Trauma, 8(4): 322-7 (Aug 1994).

AUTHORS:

1. D. Venkateswara Rao
2. G. Chaitanya
3. Anvesh Sangepu

PARTICULARS OF CONTRIBUTORS:

1. Associate Professor, Department of Orthopaedics, SMC/GGH, Vijayawada.
2. Post Graduate, Department of Orthopaedics, SMC/GGH, Vijayawada.
3. Post Graduate, Department of Orthopaedics, SMC/GGH, Vijayawada.

FINANCIAL OR OTHER

COMPETING INTERESTS: None

NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:

Dr. D. Venkateswara Rao,
M. S. (Ortho), M. Ch. (Ortho),
4th Lane, Subbarao Colony,
Flat No. 76, H. No. 54/20/2-7A,
Opp. Chaitanya College Ladies Hostel,
Near Gurudwar Temple,
Gurunanak Colony, Vijayawada-520008.
E-mail: d_yenkee@yahoo.com

Date of Submission: 08/07/2015.
Date of Peer Review: 09/07/2015.
Date of Acceptance: 06/08/2015.
Date of Publishing: 12/08/2015.