

COMPARATIVE STUDY OF QUALITY OF LIFE AMONG PATIENTS WITH COPD, BRONCHIAL ASTHMA AND HEALTHY INDIVIDUALSManju Bhaskar¹, Mayank Sarawag², Sameer Saharan³, Rajinderpal Singh⁴, Kuldeep Singh Yadav⁵**HOW TO CITE THIS ARTICLE:**

Manju Bhaskar, Mayank Sarawag, Sameer Saharan, Rajinderpal Singh, Kuldeep Singh Yadav. "Comparative Study of Quality of Life among Patients with COPD, Bronchial Asthma and Healthy Individuals". Journal of Evolution of Medical and Dental Sciences 2014; Vol. 3, Issue 54, October 20; Page: 12429-12437, DOI: 10.14260/jemds/2014/3645

ABSTRACT: The present study was conducted to compare quality of life of patients with Chronic obstructive pulmonary disease (COPD), bronchial asthma and healthy controls in different domains namely, physical health and mental health components, in addition overall quality of life and general health hence, a sample of 90 patients was used, in which 30 were COPD patients, 30 were bronchial asthma patients and 30 were healthy controls. Data was analyzed by using T-test, Chi-square test, Analysis of variance (ANOVA), and Bonferroni multiple comparisons. The findings revealed that adult male patients with COPD have poor quality of life as compared to bronchial asthmatics patients and healthy controls. The results were exhaustively discussed in the light of existing findings and other possible explanations of the findings were also offered.

KEYWORDS: Quality of life, COPD, bronchial asthma.

INTRODUCTION: Quality of life is the degree of well-being felt by an individual or group of people. It consists of two components: physical and psychological. The physical aspect includes such things as health, diet, and protection against pain and disease. The psychological aspect includes worry, stress, pleasure and other positive or negative emotional states. WHO defines quality of life as "a state dependent on physical and functional status and the degree of family support, social activity and friendship, personal achievement and philosophy; and financial adequacy and work achievement?"

The Center for Disease Control and Prevention defined quality of life as the perception of physical and mental health over time. Quality of life has assumed a special importance in the medical field in the wake of progressive move towards re humanizing high tech medicine. Quality of life is clearly becoming an important part of clinical decision making both in individual patient care and in clinical trials for the purpose of evaluating and deciding among treatment alternative. There are many states of life that are worse than death, and hence it is difficult for many to accept the "Life at any cost" principle.

Quality of life measure serve as a common denominator for comparing the overall impact of different health interventions, both at the individual and community needs. Illness and its treatment can have a major impact on many aspects that are highly relevant to the individual quality of life, such as cognitive, emotional and sexual functioning, life satisfaction and the ability to fulfill economic and other social roles. Many quality of life measures have been developed, some of which measure quality of life from a specific perspective and others, which cover a general, wider perspective.

While generic measures are broadly applicable across, types and severities of disease, across different medical treatments and across demographic and cultural subgroups, specific measures assess specific diagnostic group of patient population often with the goal of measuring responsiveness or clinically important changes. The performance of basic social roles and activities of daily living (ADL) is often used as a standard for the impact of a disease entity on quality of life.

ORIGINAL ARTICLE

A relatively early study was conducted by Barstow (1974). She reported that “major changes were evident in the style of living” manifested by the COPD patients in her study. These included alterations in bathing, grooming, dressing, eating, sleeping and mobility. Norwood (2007) reported that smoking, COPD and depression are inter-related in a sort of trinity, with depression playing a role in the initiation and maintenance of smoking, smoking leading to the development of COPD and COPD, in turn contributing to the genesis of depression.¹

Another case control study on elderly patients was conducted by Yohannes et al in which 96 Subjects were elderly outpatients with irreversible COPD aged between 70-93 years. They found that sample with epilepsy had better health related quality of life score than other chronic diseases.² Stavem et al (2000) studied the association of health related quality of life is associated with arterial PaO₂ in chronic obstructive pulmonary disease.³ Yen et al (2000) found association between PaO₂ and quality of scores in moderately to severely affected COPD patients was moderate, but higher than previously reported.⁴

This study was conducted to examine the association of respiratory symptoms and COPD severity with HRQoL (health related quality of life). This study concluded that a high level of perceived neighborhood problems were associated with poorer QOL, poorer physical functioning, and increased depressive symptoms among people with asthma when disease severity and socio-demographic factors were taken into consideration.

Over the past ten years one of the most intensely researched and discussed topic in the field of mental health has been quality of life of persons with medical diseases³. Quality of life has great significance and importance for the person’s social, psychological and emotional as well as personal life. Since quality of life is dependent to a great extent on our physical health, psychological functioning, social relationships and environmental factors. Therefore, it is logical to assume that persons with medical diseases are likely to have relatively poor quality of life as compared to the healthy individuals.

The present study is basically designed to test this assumption. Additionally it would also be interesting to compare COPD and bronchial asthmatic patients with healthy individuals in different domains of quality of life. Thus the present study also addressed this issue.

AIMS & OBJECTIVES:

1. To evaluate the quality of life in patients with COPD, Bronchial Asthma and healthy individuals.
2. To compare the quality of life in patients with COPD, bronchial asthma and healthy controls.

MATERIALS AND METHODS:

PARTICIPANT: The clinical study was conducted in Father Muller Medical College, Kankanady, Mangalore, which is a multi-specialty hospital. All patients attending the outpatient and inpatient facilities of the department of Medicine with a clinical diagnosis of chronic obstructive pulmonary disease constituted the population for the study.

The study was conducted from 1st September 2008 to the 31st of August 2010. The sample for the study consisted of ninety (N=90) males of which 30 consecutive patients with chronic COPD, 30 with bronchial asthma and 30 healthy controls which were first degree non-affected relatives of COPD patients and bronchial asthma.

ORIGINAL ARTICLE

Inclusion Criteria:

- Patients with clinical diagnosis of COPD according to GOLD's criteria.⁵
- Male patients (COPD more common in males).
- Age group between 18 and 50 years.

Exclusion Criteria:

- Patients with family history or past history of psychiatric illness not attributable to COPD.
- Patients with COPD having other medical disorders like DM, Hypertension, thyroid and other endocrine disorders, renal failure and other chronic debilitating medical conditions known to cause cognitive impairment and psychiatric morbidity.
- Patients with substance dependence other than nicotine.
- Patients who refused to give consent.

PROCEDURE: This study has been cleared by the institutional ethical committee. A written informed consent was obtained from all participants both in COPD patients and control groups. The socio demographic and clinical variables were recorded in a specific proforma prepared for this clinical study.

All the participants underwent a thorough clinical examination to rule out psychopathology and medical disorders if any. Quality of life was assessed by using Short Form 36 health survey (SF-36)^{6,7} The results obtained were analyzed using T-test, Chi-square test, Analysis of variance (ANOVA), and Bonferroni multiple comparison.

RESULTS: The three samples do not significantly differ in terms of age, marital status, religion, domicile distribution, education, occupation and income. This fact indicates that the chronic obstructive pulmonary disease (COPD) patients and the two control groups are matched.

| | Group | N | Mean | Std. Deviation | Std. Error | 95% Confidence Interval for Mean | | ANOVA F | P |
|---------------------------|------------------|-----------|---------------|----------------|--------------|----------------------------------|---------------|---------|--------|
| | | | | | | Lower Bound | Upper Bound | | |
| PHYSICAL FUNCTIONING (PF) | COPD | 30 | 85.000 | 13.896 | 2.537 | 79.811 | 90.189 | 18.664 | <0.001 |
| | Bronchial Asthma | 30 | 94.167 | 7.887 | 1.440 | 91.221 | 97.112 | | |
| | Healthy Controls | 30 | 99.500 | 2.013 | .368 | 98.748 | 100.252 | | |
| | Total | 90 | 92.889 | 10.990 | 1.158 | 90.587 | 95.191 | | |
| ROLE-PHYSICAL (RP) | COPD | 30 | 71.167 | 32.654 | 5.962 | 58.974 | 83.360 | 15.243 | <0.001 |
| | Bronchial Asthma | 30 | 95.000 | 13.772 | 2.514 | 89.858 | 100.142 | | |
| | Healthy Controls | 30 | 98.333 | 6.343 | 1.158 | 95.965 | 100.702 | | |
| | Total | 90 | 88.167 | 23.882 | 2.517 | 83.165 | 93.169 | | |

ORIGINAL ARTICLE

| | | | | | | | | | |
|----------------------------|------------------|-----------|---------------|---------------|--------------|---------------|---------------|--------|--------|
| ROLE EMOTIONAL (RF) | COPD | 30 | 74.444 | 31.176 | 5.692 | 62.803 | 86.086 | 8.003 | .001 |
| | Bronchial Asthma | 30 | 87.774 | 22.295 | 4.070 | 79.449 | 96.099 | | |
| | Healthy Controls | 30 | 97.775 | 8.466 | 1.546 | 94.619 | 100.937 | | |
| | Total | 90 | 86.665 | 24.380 | 2.570 | 81.558 | 91.771 | | |
| VITALITY (VT) | COPD | 30 | 62.667 | 31.038 | 5.667 | 51.077 | 74.256 | 17.160 | <0.001 |
| | Bronchial Asthma | 30 | 85.667 | 17.056 | 3.114 | 79.298 | 92.036 | | |
| | Healthy Controls | 30 | 93.333 | 9.034 | 1.649 | 89.960 | 96.707 | | |
| | Total | 90 | 80.556 | 24.637 | 2.597 | 75.395 | 85.716 | | |
| MENTAL HEALTH (MH) | COPD | 30 | 72.533 | 24.017 | 4.385 | 63.565 | 81.501 | 11.613 | <0.001 |
| | Bronchial Asthma | 30 | 87.867 | 11.434 | 2.088 | 83.597 | 92.136 | | |
| | Healthy Controls | 30 | 91.567 | 9.043 | 1.651 | 88.190 | 94.943 | | |
| | Total | 90 | 83.989 | 18.051 | 1.903 | 80.208 | 87.770 | | |
| SOCIAL FUNCTIONING (SF) | COPD | 30 | 72.333 | 22.504 | 4.109 | 63.930 | 80.737 | 18.753 | <0.001 |
| | bronchial asthma | 30 | 90.833 | 11.340 | 2.070 | 86.599 | 95.068 | | |
| | Healthy controls | 30 | 94.667 | 7.032 | 1.284 | 92.041 | 97.293 | | |
| | Total | 90 | 85.944 | 17.866 | 1.883 | 82.203 | 89.686 | | |
| BODILY PAIN (BP) | COPD | 30 | 73.167 | 20.171 | 3.683 | 65.635 | 80.699 | 18.083 | <0.001 |
| | Bronchial Asthma | 30 | 88.417 | 13.732 | 2.507 | 83.289 | 93.544 | | |
| | Healthy controls | 30 | 95.750 | 8.072 | 1.474 | 92.736 | 98.764 | | |
| | Total | 90 | 85.778 | 17.456 | 1.840 | 82.122 | 89.434 | | |
| GENERAL HEALTH (GH) | COPD | 30 | 61.667 | 32.625 | 5.956 | 49.484 | 73.849 | 17.800 | <0.001 |
| | Bronchial Asthma | 30 | 86.167 | 18.132 | 3.310 | 79.396 | 92.937 | | |
| | Healthy Controls | 30 | 94.667 | 9.553 | 1.744 | 91.099 | 98.234 | | |
| | Total | 90 | 80.833 | 26.107 | 2.752 | 75.365 | 86.301 | | |
| PHYSICAL COMPONENT SUMMARY | COPD | 30 | 73.483 | 21.878 | 3.994 | 65.313 | 81.652 | 22.208 | <0.001 |
| | Bronchial Asthma | 30 | 90.938 | 10.493 | 1.916 | 87.020 | 94.856 | | |
| | Healthy Controls | 30 | 97.062 | 4.228 | .772 | 95.483 | 98.640 | | |
| | Total | 90 | 87.161 | 17.280 | 1.821 | 83.542 | 90.780 | | |

ORIGINAL ARTICLE

| | | | | | | | | | |
|--------------------------|------------------|-----------|---------------|---------------|--------------|---------------|---------------|--------|--------|
| MENTAL COMPONENT SUMMARY | COPD | 30 | 69.739 | 23.826 | 4.350 | 60.842 | 78.636 | 18.081 | <0.001 |
| | Bronchial Asthma | 30 | 88.061 | 13.733 | 2.507 | 82.933 | 93.188 | | |
| | Healthy Controls | 30 | 94.240 | 7.205 | 1.315 | 91.550 | 96.931 | | |
| | Total | 90 | 84.013 | 19.308 | 2.035 | 79.969 | 88.057 | | |

Table 1: Short Form Survey 36

Table 1: comparison of all three groups (COPD, BA, Healthy Controls) on different domains of SF-36 which were Physical Functioning, Role Physical, Role Emotional, Vitality, Mental Health, Social Functioning, Bodily Pain, General Health, Physical Component Summary, Mental Component Summary. Results of SF-36 revealed that quality of life among all the three groups, COPD patients have lowest quality of life followed by Bronchial Asthma patients. Healthy controls have high QOL in all the domains of SF-36. When statistical analysis done, it shows highly significant difference in all the domains of SF-36 ($P < 0.001$), among the three groups (Table 1).

| Dependent Variable | (I) Group | (J) Group | Mean Difference (I-J) | Std. Error | P value |
|---------------------------|------------------|------------------|-----------------------|------------|---------|
| PHYSICAL FUNCTIONING (PF) | COPD | Bronchial Asthma | -9.167 | 2.401 | .001 |
| | | Healthy Controls | -14.500 | 2.401 | <0.001 |
| | Bronchial Asthma | Healthy Controls | -5.333 | 2.401 | .087 |
| ROLE-PHYSICAL (RP) | COPD | Bronchial Asthma | -23.833 | 5.367 | <0.001 |
| | | Healthy Controls | -27.167 | 5.367 | <0.001 |
| | Bronchial Asthma | Healthy Controls | -3.333 | 5.367 | 1.000 |
| ROLE EMOTIONAL (RF) | COPD | bronchial asthma | -13.330 | 5.851 | .076 |
| | | Healthy Controls | -23.331 | 5.851 | <0.001 |
| | Bronchial Asthma | Healthy Controls | -10.001 | 5.851 | .273 |
| VITALITY (VT) | COPD | Bronchial Asthma | -23.000 | 5.448 | <0.001 |
| | | Healthy Controls | -30.667 | 5.448 | <0.001 |
| | Bronchial Asthma | Healthy Controls | -7.667 | 5.448 | .489 |
| MENTAL HEALTH (MH) | COPD | Bronchial Asthma | -15.333 | 4.188 | .001 |
| | | Healthy Controls | -19.033 | 4.188 | <0.001 |

ORIGINAL ARTICLE

| | | | | | |
|----------------------------|------------------|------------------|---------|-------|--------|
| | Bronchial Asthma | Healthy Controls | -3.700 | 4.188 | 1.000 |
| SOCIAL FUNCTIONING (SF) | COPD | Bronchial Asthma | -18.500 | 3.900 | <0.001 |
| | | Healthy Controls | -22.333 | 3.900 | <0.001 |
| | Bronchial Asthma | Healthy Controls | -3.833 | 3.900 | .985 |
| BODILY PAIN (BP) | COPD | Bronchial Asthma | -15.250 | 3.831 | <0.001 |
| | | Healthy Controls | -22.583 | 3.831 | <0.001 |
| | Bronchial Asthma | Healthy Controls | -7.333 | 3.831 | .177 |
| GENERAL HEALTH (GH) | COPD | Bronchial Asthma | -24.500 | 5.743 | <0.001 |
| | | Healthy Controls | -33.000 | 5.743 | <0.001 |
| | Bronchial Asthma | Healthy Controls | -8.500 | 5.743 | .427 |
| PHYSICAL COMPONENT SUMMARY | COPD | Bronchial Asthma | -17.455 | 3.672 | <0.001 |
| | | Healthy Controls | -23.579 | 3.672 | <0.001 |
| | Bronchial Asthma | Healthy Controls | -6.124 | 3.672 | .297 |
| MENTAL COMPONENT SUMMARY | COPD | Bronchial Asthma | -18.322 | 4.238 | <0.001 |
| | | Healthy Controls | -24.501 | 4.238 | <0.001 |
| | Bronchial Asthma | Healthy Controls | -6.180 | 4.238 | .445 |

Table 2: Multiple Comparisons

TABLE 2 shows there is highly significant difference between COPD and bronchial asthma patients in all the domains of SF-36 ($p < 0.001$) except in the domain of role emotional in which there is no difference between these two groups. There is highly significant difference in all the domains of SF-36 ($p < 0.001$) between COPD patients and healthy controls. There is no statistically significant difference between bronchial asthma patients and healthy controls in the domains of SF-36.

DISCUSSION: COPD is a chronic condition with numerous psychological and physical consequences. Quality of life is generally found to be poor in COPD and other chronic pulmonary diseases. Similar findings found by Mohammed A. Zamzam et al (2012) that quality of life is impaired in patients with COPD and it deteriorates considerably with increasing severity of disease.⁸ Quality of life is worsened by co-morbid psychopathology and cognitive impairment.

ORIGINAL ARTICLE

Present investigation reveals that patients with COPD have poor quality of life both in mental component summary and physical component summary of SF – 36. These findings are consistent with those of earlier investigations. Earlier studies by McSweeney et al (1982),⁹ Prigatano et al (1984),¹⁰ Williams SJ, Burry MR (1989)¹¹ and Schrier AC (1990)¹² have used profile of mood states, sickness impact profile, Katz adjustment scale, respiratory quality of life in COPD patients and all these studies report that in COPD patients ambulation, mobility, a variety of recreational activities, sleep, rest, physical functioning, emotional functioning all are affected.

They have major problems in areas including difficulty in breathing on day to day activities and fatigue. Studies done by Yen et al (2006)⁴ and Goodridge et al (2009)¹³ using SF – 36 and SF – 12 to measure quality life reports similar findings.

Present study reveals that quality of life is significantly related to the severity of COPD. Similar finding is reported in an earlier studies, Goodridge et al (2009)¹³ Ferrer et al (1997).¹⁴ Present study also indicates that quality of life is poor in COPD patients compared to Bronchial asthma patients which are consistent with an earlier study Mark C (2011),¹⁵ using SF – 36. Present study finds that quality of life in patients with bronchial asthma is poorer in comparison to healthy controls.

Present investigation reveals that quality of life is significantly related to steroid use in COPD and bronchial asthma patients. Study done by Weldam et al. (2013) indicated that in COPD patients, the combination of illness perceptions and depressive symptoms contribute to HRQoL. More positive illness perceptions about COPD and lower levels of depressive symptoms were associated with better HRQoL.¹⁶

MERITS & LIMITATIONS: The population of this study is a selected one which does not have the characteristics of the general population of patients with COPD. Hence the samples as well as the controls are not representative of the general population. Recruitment of consecutive patients ensures that there is no sample bias. The inclusion and exclusion criteria are specific. Hence the sample consists of homogenous group of COPD patients who are otherwise not compromised. But a larger sample size will be required to enhance the reliability and validity of the results.

The present study is a comparative cross sectional case control study examining the quality of life in COPD, Bronchial asthma patients and healthy individuals. The subjects are assessed on one occasion only. The tools used have adequate established reliability and validity. All the tools are rater friendly, easy to administer, less time consuming thereby causing no discomfort to the patients. The assessment is not blind due to constraints of the study, therefore rater bias is possible.

Absence of sample selection bias, homogenous uncompromised sample of COPD patients and the control groups which includes bronchial asthma patients are conspicuous merits of this study. This is one of the few studies which compared COPD patients with bronchial asthma patients in terms of quality of life. Despite its limitations the present study definitely indicates that there is significant lower quality of life in patients with COPD in comparison to bronchial asthma patients and healthy individuals.

CONCLUSION: Present study concludes that adult male patients with chronic obstructive pulmonary disease (COPD) have poor quality of life as compared to bronchial asthma patients and healthy controls.

ORIGINAL ARTICLE

SUGGESTIONS FOR FUTURE STUDIES: Further research should ideally address the following issues:

Selection of samples which are representative of general population, Larger sample size, Use of more appropriate tools for assessment of quality of life, Blind assessment and Prospective assessment on multiple occasions.

REFERENCES:

1. Norwood RJ. A review of etiologies of depression in COPD. *International Journal of COPD* 2007; 2(4): 485-491.
2. Yohannes MA, Jamal R, Robert CB, Martin JC Depression in elderly outpatients with disabling chronic obstructive pulmonary disease. *Age and Ageing*. 1997; 27 (2):155-160.
3. Stavem K, Lossius MI, Kvein TK, et al. The health related quality of life of patients with epilepsy compared with angina pectoris, rheumatoid arthritis, asthma and chronic obstructive pulmonary disease. *Quality of life research*.2000; 9: 865-871.
4. Yen IH, Yelin EH, Katz P, Eisner MD, Blanc PD. Perceived neighborhood problems and Quality of Life, Physical Functioning, and Depressive Symptoms among adults with Asthma. *Am J Public Health*. 2006; 96: 873–879.
5. Reilly JJ, Silverman EK, Shapiro SD. chronic obstructive pulmonary disease: In Harrison's Principles of Internal Medicine., eds-Kasper DL, Braunwald E, Fauci AS, Hauser SL, Longo DL, Jameson JL. The McGraw Hill Companies. 2005; pp: 1547-1553.
6. Ware JE, Sherbourne CD. The MOS- 36 item short form health survey (SF-36), conceptual framework and item selection *medical care* 1992; 30: 473-483.
7. Scott KM, Tobias MI, Sarfati D, Haslett S. SF -36 Health survey reliability, validity and norms for New Zealand. *Australia and New Zealand Journal of Public Health* 1999; 23: 401-406.
8. Mahammed A. Zamzam, Nourane Y.Azad, Rabab A EI Wahsh, Afaf Z. Ragab, Enas M, Alam. Quality of life in COPD patients. *Egyptian Journal of Chest Diseases and Tuberculosis* 2012; 61: 281–289
9. McSweeney AJ, Grant I, Heaton RK et al. Life quality of patients with chronic obstructive pulmonary disease. *Archives of Internal Medicine*. 1982; 142:472-478.
10. Prigatano G.P, Wright EC, Levin D. Quality of life and its predictors in patients with mild hypoxemia and chronic obstructive pulmonary disease. *Archives of Internal Medicine*.1984; 144: 1613-1619.
11. Williams SJ, Burry MR. Impairment, disability and handicap in chronic respiratory illness. *Social Science and Medicine*.1989; 29: 609-616.
12. Schrier AC, Decker FW, Kaptein AA. Quality of life in elderly patients with chronic nonspecific lung disease seen in family practice. *Chest*.1990; 98: 894-899.
13. Goodridge D, Duggleby W, Gjerve J, Rennie D. Exploring the quality of dying of patients with chronic obstructive pulmonary disease in the intensive care unit: a mixed methods study; *nursing in critical care* 2009;14 (2):51-60.
14. Ferrer M, Alonso J, Morera J, et el. Chronic obstructive pulmonary disease stage and health-related quality of life. The quality of life of chronic obstructive pulmonary disease group. *Annals of Internal Medicine*.1997; 127: 1072-1079.
15. Mark C, COPD significantly reduces health-related quality of life, *Respir. Med*. 2011; 105:57-66.

ORIGINAL ARTICLE

16. Saskia WM Weldam, Jan-Willem J Lammers, Rogier L Decates, Marieke J Schuurmans. Daily activities and health-related quality of life in patients with chronic obstructive pulmonary disease: psychological determinants: a cross-sectional study. *Health and Quality of Life Outcomes* 2013; 11:190.

AUTHORS:

1. Manju Bhaskar
2. Mayank Sarawag
3. Sameer Saharan
4. Rajinderpal Singh
5. Kuldeep Singh Yadav

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Psychiatry, Mahatma Gandhi Medical College and Hospital, Jaipur, Rajasthan.
2. Assistant Professor, Department of Medicine, Jhalawar Medical College and Hospital, Jhalawar, Rajasthan.
3. Resident, Department of Psychiatry, Mahatma Gandhi Medical College and Hospital, Jaipur.

4. Resident, Department of Psychiatry, Mahatma Gandhi Medical College and Hospital, Jaipur.
5. Resident, Department of Psychiatry, Mahatma Gandhi Medical College and Hospital, Jaipur.

NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Manju Bhaskar,
#12, Jaishree Nagar,
Behind Fortis Hospital,
Malviya Nagar, Jaipur, Rajasthan.
Email: manjubhaskar123@gmail.com

Date of Submission: 08/09/2014.
Date of Peer Review: 09/09/2014.
Date of Acceptance: 15/10/2014.
Date of Publishing: 17/10/2014.