

## EVALUATION OF MODIFIED ALVARADO SCORE IN PREOPERATIVE DIAGNOSIS OF ACUTE APPENDICITIS

Ramachandra J<sup>1</sup>, Sudhir M<sup>2</sup>, Sathyanarayana B.A<sup>3</sup>

### HOW TO CITE THIS ARTICLE:

Ramachandra J, Sudhir M, Sathyanarayana BA. "Evaluation of modified alvarado score in preoperative diagnosis of acute appendicitis". Journal of Evolution of Medical and Dental Sciences 2013; Vol. 2, Issue 46, November 18; Page: 9019-9029.

### ABSTRACT:

**BACKGROUND AND OBJECTIVES:** Decision making in case of acute appendicitis may be difficult, especially for junior surgeon. Radiological investigations do not appear to be helpful sometime. A decision to operate based on clinical suspicion alone can lead to removal of normal appendix in 15-30% cases. In some studies Modified Alvarado Scoring System (MASS) was helpful in minimizing unnecessary appendectomies. The present study aims to evaluate the efficiency of Modified Alvarado scoring in preoperative diagnosis of acute appendicitis.

**METHODS:** A 100 consecutive patients suspected of acute appendicitis who were admitted, investigated and treated were taken for the study. They were prospectively evaluated using the modified Alvarado scoring (MAS) to determine whether or not they had acute appendicitis they were assigned in three groups they were treated accordingly. The MAS was correlated with operative and histopathological findings.

**RESULTS:** The results of the study showed that high score in men and children (7-9) had a sensitivity of 95.45% and 87.50% respectively, where in females it had a sensitivity of 76%. The score (5-6) in men and females had a sensitivity of 57.14% and 75% respectively.

**INTERPRETATION AND CONCLUSION:** the high score in Modified Alvarado Score Is dependable aid in the early diagnosis of acute appendicitis in men and children but is a less dependable aid as far as women are concerned.

**KEY WORDS:** Acute appendicitis, Modified Alvarado Score.

**INTRODUCTION:** One of the commonest clinical presentation that requires emergency surgery is acute appendicitis<sup>1,2</sup>. It is rare in infancy and amongst the elderly, but is common in children, teenagers and young adults<sup>3</sup>. Much efforts have been directed towards early diagnosis and intervention as approximately 6% of the population will suffer from this disease during their life time<sup>4</sup>. Delay in diagnosis leads to increase morbidity and costs.

Despite attempts to increase the diagnostic accuracy in cases of acute appendicitis, the rate of misdiagnosis in developed countries has remained constant at 15.3%<sup>5</sup>.

The classical signs and symptoms of acute appendicitis were first reported by Fitz in 1886. Since then it has remained the most common diagnosis for hospital admission requiring laparotomy<sup>1,2</sup>.

A negative appendicectomy rate of 20% has been described in the surgical literature.

# ORIGINAL ARTICLE

The diagnosis of appendicitis can be difficult, occasionally taxing the diagnostic skills of even the most experienced surgeon. Attempts to increase the diagnostic accuracy of acute appendicitis have included computer aided diagnosis, imaging by ultrasonography, laparoscopy and even radioactive isotope imaging<sup>6,7,8,9</sup>. Various scoring systems have been devised to aid diagnosis. The Alvarado score was described in 1986<sup>10</sup> and has been validated in adult surgical practice. Later, was modified by Kalan et al<sup>11</sup>.

**METHODOLOGY:** In this study, over a period of 20 months (November 2008 to June 2010) 100 patients presenting with pain in the right lower quadrant of abdomen, lasting fewer than 7 days who after clinical examination were provisionally diagnosed to have acute appendicitis and warranting surgery for the same were evaluated using the scoring system – Modified Alvarado Score.

The study was conducted on the patients presenting with clinical features suggestive of acute appendicitis admitted in surgical wards.

**Inclusion Criteria:** Patients with provisional clinical diagnosis of acute appendicitis.

**Exclusion Criteria:**

1. Patients with generalized peritonitis due to appendicular perforation.
2. Patients with appendicular mass or abscess.

**Collection of Data:** A total of 100 consecutive cases of suspected acute appendicitis who were admitted, investigated and treated were taken for the study. After detailed examination and investigations a modified Alvarado score was applied to each case.

The diagnostic scoring systems have been developed in an attempt to improve the diagnostic accuracy of acute appendicitis<sup>39</sup>. The most prominent of those scores is modified Alvarado score.

**Modified Alvarado Score:** This consists of 3-symptoms, 3 signs and a laboratory finding as described by Alvarado and later modified by Kalan et al.

SYMPTOMS/ SIGNS/ INVESTIGATIONS	SCORE	
	Yes	No
<b>Symptoms</b>		
Migration of pain to right iliac fossa	1	--
Anorexia	1	--
Nausea/ vomiting	1	--
<b>Signs</b>		
Tenderness over right iliac fossa	2	--
Rebound tenderness over right iliac fossa	1	--
Temperature >37.3°C	1	--
<b>Investigations</b>		
Leucocytosis >10 x 10 <sup>9</sup> /L	2	--
<b>Total Score</b>	<b>9</b>	<b>--</b>

Table-1: Symptoms/ Signs/ Investigation

# ORIGINAL ARTICLE

## Following decisions were taken:

Cases with score of 1-4 were observed and not operated and were followed up after discharge for next 6 months for development of acute appendicitis.

Cases with score of 5-6 were observed for next 24 hours for revision of scoring. If scores become  $\geq 7$  or their clinical condition was highly suspicious of acute appendicitis as decided by treating surgeon they were subjected for appendicectomy.

All patients who were considered for appendicectomy underwent ultrasonography of abdomen primarily to rule out other conditions mimicking acute appendicitis.

Patients with score of 7-9 who were considered candidates for appendicectomy were assessed again after ultrasonography. If any other conditions mimicking acute appendicitis were found in them. They were not operated and were considered as false positive cases.

All the specimens of appendix were sent for histopathological confirmation of acute appendicitis. Final correlation between the scoring system and final diagnosis was made.

**RESULTS:** The various features of the study which included age and sex of patient, clinical presentation, operative findings, histopathological examination were observed and analyzed. Statistical analysis of these observations and results of the study was done and presented in tabular form.

### 1. Sex Distribution:

SEX	No. of patients	Percentage (%)
Male	64	64%
Female	36	36%
<b>Total</b>	<b>100</b>	<b>100%</b>

Table No. 2: Sex Distribution

In this study number of male patients (64) were more than the number of female patients (36).

### 2. Age Distribution:

Age group (years)	No. of patients	Percentage
0 - 10	07	07
11 - 20	39	39
21 - 30	32	32
31 - 40	15	15
41 - 50	04	04
51 - 60	01	01
61 - 70	02	02
<b>Total</b>	<b>100</b>	<b>100.00</b>

Table No.3: Age Distribution

# ORIGINAL ARTICLE

Out of 100 patients, maximum patients were in the age group of 11-20 years – 39%. Next, maximum patients were in the age group of 21-30 years -32%. Only 2% were in the age group of 61-70 years, 1% in age group of 51-60 years, 4% in 41-50 years, 15% in the age group of 31-40 years and 7% in age group of 0-10 years.

**3. Division of Patient According To Score:** Patients were divided into three groups according to Modified Alvarado score as follows, and the results compared with the operative and histopathological findings.

Group-I (Score 1-4)	Conservative treatment.
Group-II (Score 5-6)	Re-assessed after few hours. Those settling were discharged, while those deteriorating with increasing scores were operated.
Group-III (Score 7-9)	Operative treatment.

GROUP	No. of patients	Percentage
Group-I	22	22
Group-II	34	34
Group-III	44	44
Total	100	100.00

**Table- 4: Division of Patient According To Score**

The above observations were made, in group-I number of patients were 22, in group-II there were 34 and in group-III there were 44 patients.

#### 4. Clinical Features:

	No. of patients	Percentage
<b>Symptoms</b>		
Migration of pain to RIF	87	87
Anorexia	78	78
Nausea/Vomiting	74	74
<b>Signs</b>		
Tenderness over RIF	83	83
Rebound tenderness over RIF	53	53
Increased temperature	67	67
<b>Laboratory Investigations</b>		
Leucocytosis	60	60

**Table-5: Clinical Features**

Among 100 patients, 87 (87%) had migration of pain to right iliac fossa, 78 (78%) had anorexia, 74 (74%) patients had nausea/vomiting, 83 (83%) patients had tenderness over right iliac fossa, 53

# ORIGINAL ARTICLE

(53%) patients had rebound tenderness over right iliac fossa, 67 (67%) patients had fever and leucocytosis was seen in 60 patients (60%).

## 5. Ultrasonography Findings:

GROUP	USG		HPR appendicitis Positive (Histopathology Record)
	Positive	Negative	
Group-II (34)	12	22	7
Group-III (44)	36	8	39

**Table- 6: Ultrasonography Findings**

USG finding in group-II among 34, 12 patients showed appendicitis and other 22 had no evidence of appendicitis. In group-III among 44 patients, 36 had appendicitis findings on USG and 8 had normal USG findings.

## 6. Distribution of Cases According To Modified Alvarado Score

### 6.1 Distribution of Cases According To Modified Alvarado Score (5-6)

Category of cases	No. of cases operated	No. of cases with HPR appendicitis	No. of cases without HPR appendicitis	Proportion of true positive
Male (n=22)	7	4	3	57.14
Female (n=10)	4	3	1	75.00
Child (n=2)	0	0	0	0
Total (n=34)	11	7	4	63.63

**Table-7.1 : Distribution of Cases According To Modified Alvarado Score (5-6)**

4 out of 7 males and 3 out of 4 females had acute appendicitis. The overall negative appendicectomy rate of patients with scores  $\leq 6$  was 36.37%.

### 6.2: Distribution of Cases According To Modified Alvarado Score (7-9)

Category of cases	No. of cases operated	No. of cases with HPR appendicitis	No. of cases without HPR appendicitis	Proportion of true positive
Male (n=22)	22	21	1	95.45
Female (n=14)	14	11	3	78.57
Child (n=8)	8	7	1	87.50
Total (n=44)	44	39	5	88.63

**Table-7.2: Distribution of Cases According To Modified Alvarado Score (7-9)**

39 cases out of 44 cases had acute appendicitis. The sensitivity of modified Alvarado score of  $\geq 7$  was 88.63%. The sensitivity was low in females 78.57% and highest in males 95.45% and in

# ORIGINAL ARTICLE

children it was 87.50%. Negative appendectomy rate was 4.54% among males, 21.43% among females and 12.5% among children.

## 7. Correlation of Age and gender according to score

	Group I (n=22)	Group II (n=34)	Group III (n=44)	P value
<b>Age in years</b>				
• 1-20	6(27.3%)	19(55.9%)	21(47.7%)	0.016*
• 21-40	16(72.7%)	15(44.1%)	16(36.4%)	
• 41-60	0	0	5(11.4%)	
• >60	0	0	2(4.5%)	
<b>Gender</b>				
• Male	13(59.1%)	24(70.6%)	27(61.4%)	0.639
• Female	9(40.9%)	10(29.4%)	17(38.6%)	

Table-8: Correlation of Age and gender according to score

## 8. Correlation of Symptoms and Signs according to score.

Symptoms and Signs	Group I (n=22)	Group II (n=34)	Group III (n=44)	P value
Migration of Pain to RIF	20 (90.9%)	27(79.4%)	40(90.9%)	0.325
Nausea/ Vomiting	12(54.5%)	20(58.8%)	39(88.6%)	0.002**
Anorexia	16(72.7%)	20(58.8%)	39(88.6%)	0.009**
Tenderness over RIF	7(31.8%)	29(85.3%)	44(100.0%)	<0.001**
Rebound tenderness	3(13.6%)	11(32.4%)	35(79.5%)	<0.001**
Temp >37.3°C	18(81.8%)	20(58.8%)	27(61.4%)	0.174
Leukocytosis	1(4.5%)	13(38.2%)	42(95.4%)	<0.001**

Table-9: Correlation of Symptoms and Signs according to score

## 9. Correlation of HPR and USG according to Total score.

Symptoms and Signs	Group I (n=22)	Group II (n=34)	Group III (n=44)	P value
HPR	0	6(17.6%)	39(88.6%)	<0.001**
USG	0	12(35.3%)	36(81.8%)	<0.001**

Table-10: Correlation of HPR and USG according to Total score

**P values are obtained by chi-square test/Fisher Exact test.**

**DISCUSSION:** Acute appendicitis being one of the commonest surgical abdominal emergencies with lifetime prevalence of approximately 1 in 7<sup>45</sup>, its diagnosis can sometimes be difficult. In an attempt to prevent negative appendectomy modified Alvarado score can be used.

## ORIGINAL ARTICLE

Hence, in the present study, an attempt is made to evaluate the efficiency of modified Alvarado score in pre-operative diagnosis of acute appendicitis done over a period of 20 months (Nov 2008 to June 2010), were included in this prospective study.

CATEGORY	Present study (%)	Hemant Nautiyal et al <sup>52</sup> (%)	Kalan M et al <sup>11</sup> (%)
Male	95.45	90	93.0
Female	78.57	100	67.0
Child	87.50	100	100.0
Total	88.63	96.6	83.7

**TABLE-11: Comparison Of Modified Alvarado Score (7-9) with other studies**

Sensitivity of appendicitis 95.45% for male in the present study with score of 7 to 9 correlates well with the figures of studies by Hemant Nautiyal et al<sup>52</sup> (90%) and with that of studies by Kalan M et al<sup>11</sup> (93%).

Sensitivity of acute appendicitis 78.57% for females in the present study with score of 7 to 9 correlates well with the figures of studies by Hemant Nautiyal et al<sup>52</sup> (100%) and but more than that of sensitivity of study by Kalan M<sup>11</sup> (67%).

Sensitivity of acute appendicitis 87.50% for children in the present study with score of 7 to 9 correlates well with the figures of study by Hemant Nautiyal et al<sup>52</sup> (100%), but has same sensitive compared to study conducted by Kalan M et al<sup>11</sup> (100%).

The overall sensitivity of acute appendicitis being 88.63% in the present study with score of 7 to 9 correlates well with the figures of studies by Kalan M et al<sup>11</sup> (83.7%) and Hemant Nautiyal et al<sup>52</sup> (96.6%).

CATEGORY	Present study (%)	Hemant Nautiyal et al <sup>52</sup> (%)	Kalan M et al <sup>11</sup> (%)
Male	57.14	93.75	67.0
Female	75.00	83.30	50.0
Child	--	0	0
Total	66.07	88.52	62.5

**TABLE-12: Comparison of Modified Alvarado Score (5-6) with other studies**

Among the score from 5 to 6, the following observations were made. The sensitivity of acute appendicitis 57.14% for males in our study is lesser than the sensitivity of studies by Kalan M et al<sup>11</sup> (67%) and Hemant Nautiyal et al<sup>52</sup> (93.75%).

And sensitivity of acute appendicitis 75% for females in the present study is high compared with the figures of studies by Kalan M et al<sup>11</sup> (50%) and Hemant Nautiyal et al<sup>43</sup> (83.3%). No Child in Group-II was operated.

The overall sensitivity of acute appendicitis being 66.07% in the present study with score of 5 to 6 correlates well with the figures of studies by Kalan M et al<sup>11</sup> (62.5%) and less compared to Hemant Nautiyal et al<sup>52</sup> (88.52%).

In another study by Mohanty Sudhir Kumar et al<sup>46</sup> was conducted on 45 patients prospectively using the modified Alvarado score. They found positive predictive value of 95.2% for males and for females, 93.3%, which was higher than our study.

# ORIGINAL ARTICLE

---

They concluded that the score was useful in distinguishing acute appendicitis from other acute abdominal conditions, thus decreasing negative appendicectomy.

In another study by Fente BG<sup>53</sup> was conducted on 128 patients retrospectively using the modified Alvarado score. They found that sensitivity of 92.93% and specificity of 92.93% were recorded in their study.

They concluded that the score is a simple, safe and cost effective aid in diagnosis of acute appendicitis and decreases negative appendicectomy rate.

In another study by Sanjot B. Kurane<sup>54</sup> was conducted on 60 patients prospectively using the modified Alvarado score. They found that modified Alvarado score has sensitivity of 78.26%, and specificity 83.78% and Ultrasonography had sensitivity of 82.61%, specificity of 89.19%.

They concluded that the score is a useful tool in clinical decision making. However additional information provided by ultrasonography improves diagnostic.

**CONCLUSION:** Though acute appendicitis is the commonest surgical abdominal emergency with a life time prevalence of approximate 1 in 733, its diagnosis can sometimes be difficult.

From the present study, it may be concluded that high scores (7-9) in modified Alvarado Score is a dependable aid in early diagnosis of acute appendicitis in men and children as compared to other studies, but same is not true as far as women are concerned.

USG abdomen is a useful tool in diagnosis of appendicitis (HPR positive) in patients with score 5 to 6 and 7 to 9.

Our study had almost similar results as of the studies, Bengezi et al<sup>46</sup>, Fente BG<sup>53</sup>, Sanjot B. Kurane<sup>54</sup> and as mentioned before Kalan M et al<sup>11</sup> and Hemant Nautiyal et al<sup>52</sup>.

**SUMMARY:** This study was conducted on 100 consecutive patients provisionally diagnosed to have acute appendicitis. Modified Alvarado Score was applied to all these patients.

- Maximum number of cases were seen in age group of 11-20 years, 37 cases (37%).
- 87 (87%) patients presented with pain in the right lower quadrant of abdomen, lasting fewer than 7 days.
- High scores (7-9) in men, children is dependable and in early diagnosis of acute appendicitis, whereas it is not so in case of females.

## BIBLIOGRAPHY:

1. Puylaart JB. Acute appendicitis ultrasound evaluation using graded compression. *Radiol.* 1986; 158: 355-60.
2. Pearson RH. Ultrasonography for diagnosing the appendicitis. *Br Med J.* 1988; 297: 309-10.
3. Jones DJ. Appendicitis. *Br Med J.* 1993; 301: 207-10.
4. Anonymous. A sound approach to the diagnosis of acute appendicitis (editorial). *Lancet.* 1987; 1: 198-200.
5. Jaffee BM, Berger DH, Brunnicardi FC, Anderson DK, Billion TR, Dunn DL (eds.). The Appendix. In: Schwartz's Principles of Surgery. 8<sup>th</sup> ed., New York: McGraw Hill; 2005. P. 1119-55.
6. Balthazar EJ, Megibow AJ et al. C of appendicitis. *Am J Radiol.* 1986; 6: 185-193.
7. Takada T et al. Ultrasonographic diagnosis of acute appendicitis in Surgical indication. *In Surg.* 1986; 71: 9-13.

## ORIGINAL ARTICLE

---

8. Clarke PJ et al. The use of laparoscopy in the management of right iliac fossa pain. *Ann R Coll Surg Engl.* 1986; 68: 68-69.
9. Eric BR et al. Tc-99-HMPAO white blood cell scan for diagnosis of acute appendicitis in patients with equivocal clinical presentation. *Ann Surg.* 1997; 226(1): 58-65.
10. Alvarado A. A practical score for the early diagnosis of acute appendicitis. *Ann Emerg Med.* 1986; 15: 557-564.
11. Kalan M et al. Evaluation of the modified Alvarado score in the diagnosis of acute appendicitis – A prospective study. *Ann R Coll Surg Engl.* 1994; 76: 418-9.
12. Fitz RH. Perforating inflammation of the vermiform appendix: With special reference to its early diagnosis and treatment. *Trans Assoc Am Physicians.* 1886; 1: 107.
13. Ellis H. Appendix. In: Schwartz SI (ed.). *Maingot's Abdominal Operations.* 8<sup>th</sup> ed. Norwalk: Appleton Century Crafts. 1985: 2: 1255.
14. McBurney C. Experience with early operative interference in cases of diseases of the vermiform appendix. *NY State Med J.* 1889; 50: 676.
15. Appendix Shackelford surgery of the alimentary tract 5<sup>th</sup> edition volume 4 chapter 13 page 180-209.
16. Saidi HS, Chovda SK. Use of modified Alvarado score in the diagnosis of acute appendicitis. *East Afr Med J.* 2003 Aug; 80(8): 411-4.
17. Lamporelli MJ, Haque HM, Pogson CJ and Ball AB. A prospective evaluation of the combined use of the modified Alvarado score with selective laparoscopy in adult females in management of suspected appendicitis. *Ann R Coll Surg Eng.* 2000 May; 82(3): 192-5.
18. Moore Persaud. The developing human. The digestive system. 7<sup>th</sup> edition. 266-273.
19. The Appendix. Chapter-5. Lee McGregor's Synopsis of Surgical Anatomy, GAG Decker, DJ du Plessis, Verghese Publishing House, Bristol. 1986; 12: 31-42.
20. Skandalakis JE, Graysu, Reckett SR: the colon and Rectum. In Skandalakis JE: Grays W (eds): *Embryology for Surgens*, Baltimore, Williams, and Wilkins 1994 : 242-281.
21. Ronan 'O' conell, "The vermiform appendix," Bailey and Love Short practice of surgery, 24<sup>th</sup> edition : 1203-1218.
22. Keith L Moore & Anne MR Agur. *Essential Clinical Anatomy.* 3<sup>rd</sup> Edn. Lippincott Williams & Wilkins. Abdomen. 155-158.
23. James M Crawford. Appendix. Chapter-17 In: Robbins & Cotran's Pathologic Basis of Diseases. Kumar, Abass, Fansto. Philadelphia, Pennsylvania, Elseviers. 2004; 7: 870-71.
24. Norman L Browse et al. Browse's Introduction to the Symptoms and Signs of Surgical Disease. 4<sup>th</sup> edn. The Abdomen : 405-406.
25. Ronan P O'Connel. The Vermiform Appendix. Chapter-67. In: Bailey & Love's Short Practice of Surgery. RCG Russell, Norman S Williams and Christopher JK, Bulsrode, London. Arnold. 25<sup>th</sup> edn. 2008: 1205-18.
26. Burkitt DP. The etiology of appendicitis. *Br J Surg.* 1971; 58: 695.
27. Butler C. Surgical pathology of acute appendicitis. *Human Patholo.* 1981; 12: 870.
28. Miranda R, Johnston AD, O'Leasy JP. Incidental appendectomy: Frequency of Pathologic Abnormalities. *Ann Surg.* 1980; 46: 355.
29. Rautio M, Soxen H, Sitonen A et al. Bacteriology of histopathologically defined appendicitis in children. *Pediatr Infect Dis J.* 2000; 19: 1078.

## ORIGINAL ARTICLE

---

30. Softer D, Zaid S, Klousner J et al. Peritoneal cultures and antibiotic treatment in patients with perforated appendicitis. *Eur J Surg*. 2001; 167: 214.
31. Charles F Brunicaudi. *Schwartz's Principles of Surgery*. 8<sup>th</sup> edn., Chapter-29. "The Appendix" : 1119-1135.
32. Wagner JM, McKinney WP, Carpenter JL. Does this patient have appendicitis? *JAMA*. 1996; 276: 1589-1594.
33. Lazar J Greenfield et al. *Surgery, scientific principles and practice*. 3<sup>rd</sup> edn., Lippincott. Chapter-52. Acute Abdomen and Appendix : 1224-35.
34. Whitney TM, Macho JR, Russell TR et al. Appendicitis in acquired immunodeficiency syndrome. *Am J Surg*. 1992; 164: 467-70.
35. Flum Dr, Steinberg SD, Sarkis AV, Wallack MK. Appendicitis in patients with acquired immunodeficiency syndrome. *J Am Coll Surg*. 1997; 184: 481-486.
36. Jose FE Fischer. *Mastery of Surgery*. 5<sup>th</sup> edition. Lippincott Chapter-129. Appendicitis & Appendiceal Abscess : 1430-33.
37. Engstrom L, Fenyo G. Appendectomy: Assessment of stump invagination versus simple ligation: A prospective randomized trial. *Br J Surg*. 1985; 72: 971-972.
38. Teicher I, Larida B, Cohen M et al. Scoring system to aid in diagnosis of appendicitis. *Ann Surg*. 1983; 198: 753-759.
39. Christian F, Christian G P, A simple scoring system to reduce the negative appendectomy rate, *Annals of the Royal College of Surgeons of England* (1992) vol. 74: 281-28.
40. Christian Ohmann, PhD; Claus Franke, MD Clinical Benefit of a Diagnostic Score for Appendicitis Results of a Prospective Interventional Study *Arch Surg*. 1999;134 :993-996.
41. Enochsson L,. Gudbjartsson T. The Fenyo" - Lindberg scoring system for appendicitis increases positive predictive value in fertile women *Surg Endosc* (2004) 18: 1509-1513.
42. Sitter H, Hoffman.S, Diagnostic score in appendicitis validation of a diagnostic score (Eskelinen score) in patients in whom acute appendicitis is suspected, *Langenbecks Arch Surg* (2004) 389:213-218.
43. Hannu Lintula, Appendicitis score for children with suspected appendicitis A randomized clinical trial, *Langenbecks Arch Surg* (2009) 394:999-1004.
44. Stephens PL, Mazzuco JJ. Comparison of ultrasound and Alvarado Score for the diagnosis of acute appendicitis: A prospective study, *Ann R Coll Surg. Eng*. 1994; 761: 418-9.
45. Mohanty Sudhir Kumar, Sal Kaushik. Evaluation of modified Alvarado Score in decreasing negative appendectomy rate – our experience, *Ind J Surg*. 2000 62(5): 342-343.
46. Bengezi, OA, Fallouji AL M. Modified Alvarado Score in Diagnosis of Acute Appendicitis, *The British Journal of Surgery*. 1997 Suppl. May; 84: 30.
47. David T Chiang<sup>1</sup>, Elaine I Tan<sup>2</sup>, David Birks To have or not to have' should computed tomography and ultrasonography be implemented as a routine workup for the workup of acute appendicitis in the regional hospital, *Ann R Coll Surg Engl* 2008;90:17-21
48. Alexandra Platon Helmi Jlassi Olivier T. Rutschmann Evaluation of a low-dose CT protocol with oral contrast for assessment of acute appendicitis, *Eur Radiol* (2009) 19: 446-454.
49. A. van Randen W. Laméris H. W. van Es Profiles of US and CT imaging features with a high probability of appendicitis, *Eur Radiol* (2010) 20: 1657-1666.

# ORIGINAL ARTICLE

---

50. Boudewijn R. Toorenvliet, Fraukje Wiersma Routine Ultrasound and Limited Computed Tomography for the Diagnosis of Acute Appendicitis, *World J Surg* (2010) 34:2278–2285.
51. Shozo Yokoyama, Katsunari Takifuji C-Reactive protein is an independent surgical indication marker for appendicitis: a retrospective study, *World Journal of Emergency Surgery* 2009, 4:36.
52. Hemant Nautiyal · Shabi Ahmad · N. K. Keshwani · D. N. Awasthi Combined use of modified Alvarado score and USG in decreasing negative appendicectomy rate *Ind J Surg.* 2010 volume 72: 42-48.
53. Fente BG. Echem RC Prospective evaluation of the Bengezi and Al-Fallouji modified Alvarado score for presumptive accurate diagnosis of acute appendicitis in University of Port Harcourt Teaching Hospital, Port Harcourt. *Nigerian Journal of Medicine*; 2009; 18(4):398-401.
54. Sanjot B. Kurane . M. S. Sangolli . A. S. Gogate A one year prospective study to compare and evaluate diagnostic accuracy of modified Alvarado score and ultrasonography in acute appendicitis, in adults *Indian J. Surg.* 2008 ; 70:125–12.
55. Shozo Yokoyama, Katsunari Takifuji C-Reactive protein is an independent surgical indication marker for appendicitis: a retrospective study, *World Journal of Emergency Surgery* 2009, 4:36.

## **AUTHORS:**

1. Ramachandra J.
2. Sudhir M.
3. Sathyanarayana B.A.

## **PARTICULARS OF CONTRIBUTORS:**

1. Professor, Department of General Surgery, Kempegowda Institute of Medical Sciences, Bangalore.
2. Associate Professor, Department of General Surgery, Kempegowda Institute of Medical Sciences, Bangalore.
3. Professor, Department of General Surgery, Kempegowda Institute of Medical Sciences, Bangalore.

## **NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:**

Dr. Ramachandra J,  
Professor, Department of General Surgery,  
Kempegowda Institute of Medical Sciences,  
V.V. Puram, Bangalore.  
Email – ramachandrasuhas@yahoo.com

Date of Submission: 30/10/2013.  
Date of Peer Review: 31/10/2013.  
Date of Acceptance: 12/11/2013.  
Date of Publishing: 15/11/2013